



FLY TIMES

ISSUE 38, April 2007

Jeffrey M. Cumming, co-editor
Invertebrate Biodiversity
Agriculture & Agri-Food Canada
CEF, Ottawa, Ontario, Canada, K1A 0C6
Tel: (613) 759-1834
FAX: (613) 759-1927
Email: cummingjm@agr.gc.ca

Art Borkent, co-editor
691 - 8th Ave SE
Salmon Arm, British Columbia
V1E 2C2, Canada
Tel: (250) 833-0931
FAX: (250) 832-2146
Email: aborkent@jetstream.net

Welcome to the latest *Fly Times*. As usual this issue contains our regular reports on meetings and activities, opportunities for dipterists, as well as information on recent and forthcoming publications.

The electronic version of the *Fly Times* continues to be hosted on the North American Dipterists Society website at <http://www.nadsdiptera.org/News/FlyTimes/Flyhome.htm>. We will, of course, continue to provide hard copies to those without web access. We would greatly appreciate your independent contributions to this newsletter. We need more reports on trips, collections, methods, etc., with associated digital images if you provide them. Feel free to make requests and to share your opinions about what is happening in your area of study, or to pass on any ideas you have on how to improve the newsletter and the website.

The *Directory of North American Dipterists* is constantly being updated and is currently available at the above website. Please check your current entry and send all corrections to Jeff Cumming.

Issue No. 39 of the *Fly Times* will appear next October. If possible, please send your contributions by email, or disc, to either co-editor. Those of you without internet access may fax, or mail hard copy contributions. All contributions for the next *Fly Times* should be in by the end of September, 2007.

NEWS

Changes at the Helm - New Editors for the Fly Times

Art and Jeff have been putting the *Fly Times* together for the past 19 years and we feel that it is time to pass the reins over to new editors, with fresh perspectives and energy. If you want to volunteer to take on the task (along with its many honours and accolades) or have suggestions or opinions on what the future of the *Fly Times* might look like, let us know.

5th Annual Meeting of the North American Black Fly Association (NABFA)

By Julio Rivera

Department of Ecology and Evolutionary Biology, University of Toronto &
Department of Natural History, Royal Ontario Museum
100 Queen's Park, M5S 2C6, Toronto ON, Canada; julior@rom.on.ca

The 5th Annual Meeting of the North American Black Fly Association was held 7-9 February 2007 at the University of Georgia in Athens, Georgia. The meeting was chaired and organized by NABFA President Jay P. Overmyer. Thirty-four workers attended the meetings and 24 presentations were given. Topics included black fly control, ecology, taxonomy, systematics and biogeography. Abstracts of these presentations will be posted in the NABFA website (<http://www.zoo.utoronto.ca/nabfa/NABFA.html>).

Among the highlights of the meeting was a guided tour of the University of Georgia's Black Fly Colony, which has been maintained continuously for more than 20 years. Another notable event was the inaugural Mike Spironello Award, presented annually to the student giving the best talk. The award was established to honour the memory of former NABFA Secretary and black fly researcher, Mike Spironello, who passed away unexpectedly in 2006. This year's winner, was Paul Smith of the University of Georgia, who delivered a co-authored presentation on the black fly vectored disease vesicular stomatitis (full title and authorship given below). Paul was presented with a copy of Adler, Currie, and Wood's monumental book *The Black Flies (Simuliidae) of North America*.

NABFA invites all black fly researchers to join NABFA and to present their research at upcoming meetings. Students are especially encouraged to participate in the Mike Spironello Award competition. Membership is free of charge. Please visit the NABFA website or contact secretary Julio Rivera (julior@rom.on.ca) for further information.

Regular Presentations:

- Reproductive Status of Cytospecies and Cytotypes in Sympatry at Three Locations in Western Montana. Gerald F. Shields, Judith A. Pickens, Lindee M. Strizich and Gregory M. Clausen. Carroll College, Helena, MT.
- The Black Fly Genome Project Proposal: Progress Report. Charles Brockhouse. Creighton University, Omaha, NE (Presented by Peter Adler).
- What's going on in Minnesota? Black Fly Monitoring and Control in the Greater Metropolitan Area of the Twin Cities of Minneapolis and St. Paul, Minnesota, USA. John Walz, Abe Benson and Carey LaMere. Metropolitan Mosquito Control District, Black Fly Control Program, St. Paul, MN.
- Phylogeny of the black fly genus *Cnephia* (Diptera: Simuliidae). Kenneth Pruess. University of Nebraska, Lincoln, NE.
- Who's in control, the fly or the fungus? Spore shape of *Harpella melusinae* in larval black flies. Charles E. Beard and Peter H. Adler. ESPS, Clemson University, Clemson, SC.
- An overview of black fly control in Hunterdon County, New Jersey. Tadhgh Rainey. Hunterdon County Department of Health, Flemington, NJ.
- Black fly vignettes. Peter H. Adler. Clemson University, Clemson, SC
- The contribution of South- and Central American lineages of black flies (Diptera: Simuliidae) to the North American fauna. Douglas C. Currie^{1,2} and Justin Ancheta². ¹Royal Ontario Museum, Toronto, ON, ²University of Toronto, Toronto ON.
- Stable isotope turnover in black fly, *Simulium vittatum* IS-7, larvae. Jay Overmyer¹, M. Aaron MacNeil² and Aaron T. Fisk³. ¹University of Georgia, Athens, GA, ²University of Newcastle, Newcastle upon Tyne, UK, ³University of Windsor, Windsor, ON.
- Effects of *Bacillus thuringiensis israelensis* (Bti) black fly treatments on non-target organisms in the Delaware River in Pennsylvania and New Jersey. Dave Rebeck. Pennsylvania Department of Environmental Protection, Division of Vector Management, Black Fly Suppression Program.
- The Vectobac 12AS story: how industry, academia, NGOs and governments collaborated to develop the premier black fly control larvicide. Ernest Dankwa. Valent Biosciences.
- The role of chromosome change in the evolution of black flies. Gerald Shields. Carroll College, Helena, MT.
- Molecular tools and the study of black fly-borne parasites. Will Reeves. Gainesville State College, Oconee Campus.
- Twenty-five years of black fly colonization, 1981-2006. Elmer Gray. University of Georgia, Athens, GA.

Poster Presentations:

- A Longitudinal Analysis of the Distributions of and an Assessment of the Reproductive Status of Two Siblings of the *Simulium arcticum* Complex at Little Prickly Pear Creek, Lewis and Clark County, Montana. Gregory M. Clausen and Gerald F. Shields. Carroll College, Helena, MT.
- Speciation in the Black Fly, *Simulium arcticum* Complex (Diptera: Simuliidae). Gerald F. Shields. Carroll College, Helena, MT.
- The *Simulium arcticum* Complex: Environmental Effects on Distribution of Taxa at Trout Creek and Reproductive Status of Taxa at the Blackfoot River. Lindee M. Strizich and Gerald F. Shields. Carroll College, Helena, MT.
- Reproductive Status of Cytotypes of the *Simulium arcticum* Complex at Rock Creek, Missoula County, Montana. Judith A. Pickens and Gerald F. Shields. Carroll College, Helena, MT.

Mike Spironello Award Competition:

- Black fly fauna of Nepal. Dustin A. Swanson and Peter H. Adler. Clemson University, Clemson, SC.
- Utility of the Cytochrome Oxidase I Gene for Species Recognition and Phylogeographic Analysis in Black Flies (Diptera: Simuliidae). Julio Rivera¹ and Douglas C. Currie^{1,2}. ¹University of Toronto, Toronto, ON, ²Royal Ontario Museum, Toronto, ON.
- Effects of temperature and developmental stage on nitrogen and carbon stable isotopes in the black fly, *Simulium vittatum* IS-7. Grant Howell and Jay Overmyer. University of Georgia, Athens.
- GAA Molecular Approach to Identifying Members of the *Simulium jenningsi* species-group. Beth Alexander. University of Tennessee, Knoxville, TN.
- Vesicular stomatitis: A research prospectus; Study of disease cycle, vector transmission, and virus maintenance. Paul Smith, Danny Mead and Ray Noblet. University of Georgia, Athens, GA.

Guest Presentation:

- Small is significant, be it organism or ecosystem. Judy Meyer. University of Georgia

**Final Announcement for the 2007 Field Meeting of the North American Dipterists Society
Silver City, New Mexico, 13-16 August 2007**

by Jim O'Hara

Invertebrate Biodiversity, Agriculture & Agri-Food Canada, 960 Carling Avenue,
Ottawa, Ontario, Canada K1A 0C6; oharaj@agr.gc.ca

The next field meeting of the North American Dipterists Society is just months away. We are expecting a good turn-out of dipterists and hope that the diverse fly fauna of the Silver City area will meet the expectations of the participants. Details about the Silver City area, including maps, pictures, and other information, were given in the First Announcement (*Fly Times* 35, October 2005) and Second Announcement (*Fly Times* 37, October 2006). This final announcement provides a few specifics about registration, accommodation, and the meeting schedule.

Registration will be held in Harlan Hall ([see PDF for location](#)) on the campus of Western New Mexico University from 3:30–5:00 p.m. and from 6:30–7:00 p.m. on Monday August 13th. The fee for registration will be US\$20/ person, payable in cash. Receipts will be issued at the time of payment. A registration fee of US\$10 will apply to accompanying persons if they plan to attend group activities such as the BBQ at the Gomez Peak Group Picnic Area on August 14th or 15th. Information about collecting in the Silver City area will be available during registration.

Persons planning to attend the NADS field meeting are asked to inform me ([Jim O'Hara](#)) of their intention by Monday June 25th so that proper local arrangements for the meeting can be made. Anyone wishing to give an oral presentation (15 minutes including questions) is asked to send me their title by June 25th. Plan to bring your presentation to the meeting on electronic media (preferably CD or memory stick). I will send out a meeting schedule, including a list of presentations, in mid July to all persons who have indicated to me that they plan to attend. I will leave for Silver City during the first week of August and will not be reachable thereafter until the meeting begins.

There will be a presentation by a guest speaker and an orientation talk by myself on the evening of Monday August 13th. The next day will consist of collecting in the morning and afternoon and oral presentations in the evening. The last two days will consist of collecting during the day and sorting/preparing specimens in the evening, unless there are so many talks planned that they have to extend into the evening of the third day. Lab space and microscopes will be available to us in Harlan Hall for the sorting of specimens. I have booked the Gomez Peak Group Picnic Area north of Silver City for our exclusive use on days two and three of the meeting (Tuesday and Wednesday). See the Second Announcement for habitats in the vicinity of the Picnic Area. Weather permitting, we will have a BBQ in the Picnic Area during the late afternoon on one of those two days. The Picnic Area is closed to use after dark.

Meals will be the responsibility of individuals during the meeting, with the exception of the BBQ at the Gomez Peak Group Picnic Area and possibly a group dinner on the evening of the last day. There are many restaurants around Silver City to choose from and a Wal-Mart with a large grocery section on Hwy. 180 east of town.

Accommodation is also the responsibility of individuals. There are a number of hotels available, mostly on Hwy. 180 east of the town center. Silver City is not very big, but the University is on the west side of town and the hotels are mostly on the east side, so the distance between them is a short car drive or a longer walk. If you are looking for reasonably-priced accommodation with an included Continental Breakfast and such amenities as a refrigerator, microwave, coffee maker, and internet service, then a good choice is the [Econo Lodge](#). We have been granted a group rate of \$69.99 per night at the Econo Lodge – call them direct at (505) 534-1111 and mention that you are a member of the North American Dipterists Society. Best to book ahead. Cheaper accommodation, more in the \$50 per night range, is available at the Motel 6 or Copper Manor Motel (Motel 8 not recommended). Check out the Internet for additional information and other options.

Trapping a Rarely Collected Fly: *Melanderomyia kahli* Kessel

by Jeff Skevington and Barry Flahey
 Canadian National Collection of Insects, Arachnids & Nematodes
 960 Carling Avenue, Ottawa, ON, K1A 0C6, Canada;
skevingtonj@agr.gc.ca; flaheyb@agr.gc.ca

Melanderomyia are rarely collected Platypezidae that feed on stinkhorn fungi as larvae. The CNC had only 1 specimen in the collection until BF managed to collect more. They are actually fairly easy to collect as might be expected given their predictable feeding habits. The first author was keen to get these flies for molecular phylogenetics analysis of the lower Cyclorrhapha and when BF arrived at the lab boasting about the big phallus in his yard, JS asked him to try to find and collect *Melanderomyia*. BF managed to collect a few in 2004 and has since perfected his technique.

A cage was placed over the stinkhorn as soon as it was detected. The sides of the cage were propped up on pieces of wood so that the completely open bottom of the cage was about 5 cm off the ground (Fig. 1). The rest of the cage was closed. BF returned to the trap at regular intervals and collected the flies that had flown up and were crawling around the top of the cage. A more passive method might be to put a cage with a collecting head at the top over the stinkhorn; however, you may end up with a bottle of calyptrates before you catch any platypezids. Not surprisingly, this technique appears to capture predominantly females.

It would be great to get more people collecting these flies. We know little about their phenology and perhaps more than one species exists. Our specimens were collected in October 2004 and October 2006 but we suspect that they occur whenever and wherever there are stinkhorns fruiting. There are some photos of platypezids on stinkhorns on BugGuide (<http://bugguide.net/node/view/19382>) that appear to be *Melanderomyia* (photographed on 21 September 2004 in North Carolina). There are oodles crawling around on stinkhorns. Based on our observations and on these photos, it appears that *Melanderomyia* may be locally common, or at least are common during a very narrow window when the stinkhorn is at its peak. Try looking late in the season if you have no luck earlier in the year. These records suggest that these may be late season flies.

Incidentally, the range of *Melanderomyia kahli* is given as Kansas, Iowa, Illinois, and Ohio on the Diptera Nomenclator (<http://www.sel.barc.usda.gov:591/diptera/names/searchno.htm>), so our collections in Ottawa represent new records for ON and Canada.



Fig. 1. Cage placed over stinkhorn fungus used for collecting *Melanderomyia kahli*. Note that the fly on the top of the cage is not a *Melanderomyia* but is a calyptrate.

The Biosystematic Database of World Diptera in 2007

by Owen Lonsdale

Entomology Department, Rm. CE-607, Smithsonian Institution
National Museum of Natural History, 10th & Constitution Ave. NW
Washington, D.C., 20560-0168, USA; neoxabea@hotmail.com

Starting this May, I am fortunate enough to begin my research as a Schlinger postdoctoral fellow in the Entomology Department of the Smithsonian Institute's Museum of Natural History. Along with Chris Thompson, I will continue to develop the Biosystematic Database of World Diptera (BDWD) and the Diptera Web Site – a comprehensive portal to knowledge about the true flies, and a framework to organize and integrate all current and future knowledge about these beautiful and highly diverse organisms. My contribution will primarily be content editing and production of the journal MYIA. I also plan to enlist the help of dipteran taxonomic specialists (such as yourselves) for peer-review, to make sure that the database meets the highest possible standards of accuracy and scientific content. As such, I strongly invite anyone developing a catalogue, taxonomic revision or faunistic review to contact us so that we may improve this resource for the dipterological community. A full description of the BDWD project can be viewed online at <http://www.sel.barc.usda.gov/Diptera/biosys.htm>, along with a link to the "Nomenclator", a powerful interactive tool that not only searches for dipteran references and names, but provides taxonomic and nomenclatural information about those names. Much of the database has already been developed through the efforts of Chris, Betty Thompson and Irina Brake (the previous Schlinger postdoctoral fellow who now works at the Natural History Museum in London), and I aim to have much more completed this coming year.

In addition to coordinating the BDWD, I plan to publish my revisionary work on the New World Clusiidae (Schizophora), which I defended this past April as my doctoral thesis. In the Americas, I currently recognize 362 species in ten genera, an amazing 220 of which I am describing for the first time – most of these belong to highly diverse *Sobarocephala*, a colourful genus of worldwide distribution endemic to the tropical Neotropics. While most of my revisionary work is yet to be published, some of my findings can be accessed online at the Tree of Life website (<http://tolweb.org/Clusiidae>). I have also posted similar pages on the families Somatiidae (<http://tolweb.org/Somatiidae>) and Acartophthalmidae (<http://tolweb.org/Acartophthalmidae>).

Surveying Dolichopodids and Damselflies in the Society Islands, French Polynesia

by Neal L. Evenhuis

Bishop Museum, 1525 Bernice Street, Honolulu,
HI 96817-0916, USA; neale@bishopmuseum.org

As part of the NSF-funded Arthropod Survey of French Polynesia, the Bishop Museum conducted an expedition to the Society Islands the last two weeks of March 2007. Ronald Englund, aquatic biologist, specializes in odonates and collected most aquatic invertebrates he encountered; I surveyed forests and aquatic habitats for dolichopodids, primarily of the genus *Campsicnemus*, which has undergone a

tremendous amount of speciation in French Polynesia with some 43 new species having been discovered prior to this expedition.

We arrived in a hot and muggy Papeete late in the evening of 17 March and got much needed sleep that night after a rather bumpy 5 1/2 hour flight down from Honolulu. This was my second trip; Ron had been to French Polynesia (FP) about 8 times already, but previous collecting was in the Marquesas, the Australs, and Raiatea (the last was pretty much a rain-out the previous year). This trip would key in on the islands of Raiatea (once again for Ron) and Huahine (first time for both of us). The expedition would start in Raiatea on Tuesday. Sunday and Monday would include trip preparation and a side trip to the highlands in central Tahiti Nui.

Early Sunday morning we caught the ferry to Moorea and after the speedy 30 minute trip between Tahiti and Moorea were met at the dock by the expedition coordinator, Elin Claridge (a weevil specialist) of the U.C. Berkeley Gump Research Station who drove us to the lab after purchasing some needed collecting supplies and *beignet de chocolat* at the local Champion store. My job that day was to sort through the last 6 months worth of Malaise trap residues (two traps on Moorea; three on Mt. Marau on Tahiti) (Fig. 1) while Elin and Ron took advantage of the good weather to collect at Vaioro Riviere on the Tohiea summit trail that Ron had not hit in his previous trips to Moorea. The Malaise trap residues turned up a few nice surprises (the robber fly *Mesoleptogaster* - a first record of the genus and family from FP; a keroplatid (*Heteropterna*), a pipunculid, and a rhagionid (*Chrysopilus*) - also firsts for FP; and a few more specimens of some new *Campsicnemus* previously represented by only one or two specimens). In just a few hours, Ron managed to secure a potential new damselfly and also a new *Campsicnemus* by sweeping rocks in the stream and adjacent seeps. When they returned from their collecting late that afternoon, we loaded up the Gump Station Land Rover with our gear for the next 12 days and drove it to the port for the ferry ride to Papeete (it would eventually be ferried from there to Raiatea and Huahine for our fieldwork on those islands).



Fig. 1. Sorting through Malaise trap residues at U.C. Berkeley Gump Research Station.

The next morning - a fair-weather one, we were met at the hotel by our good friend and botanist, Jean-Yves Meyer, deputy director for the Delegation de Recherché in FP, who drove us to a special highland bog on Tahiti called Anaorii Maracage (650 m) (Fig. 2). It is the only such high elevation bog in FP and is characterized by the endemic *Pandanus* (*P. papenooensis*). After a two-hour jeep ride up a narrow, rain-soaked, rubbly switchback road past the FP army in full camouflage (who happened to be using the road that day for a long march up the mountain), we arrived at the Maracage and hiked in through a grove of *Miconia* to a small stream that drained the bog. Here I found many specimens of a species of *Campsicnemus* that I had collected the previous trip to Tahiti that apparently restricts itself to seeps near streams; and also collected a new species of *Campsicnemus* not collected previously. We hiked into the bog, which was ankle deep in most places but knee to waist deep upon nearing the *Pandanus* that fringed the bog. After slogging through the bog and falling into waist-deep holes a few times, we made it back to our jeep and headed back down the switchbacks. Our next stop was at Lac Vaihiria (essentially a man-made reservoir slotted into the narrow valley), where we did a bit of sweep netting for flies and made some hilarious efforts at collecting the speedy *Anax* dragonflies darting along the water's edge, including (much to Jean-Yves's dismay) Ron donning snorkel and mask to "sneak up" on it from the water. After about 30 minutes, we captured one with a net for a photo op later that day back at our hotel room's jerry-rigged photo studio.



Fig. 2. Anaorii Maracage, a highland bog on Tahiti characterized by *Pandanus*.

The next morning began with a short early flight northwest from Papeete, landing on the island of Raiatea at 6:30 am where we met up with Elin and the Land Rover who had taken an overnight freighter with the truck. After a hearty breakfast and coffee at a snack shop in the main town of Uturoa, we drove to what was to be our accommodations for the next 5 days -- the Kaoha Nui Ranch Pension -- 5 km south of Uturoa past the more luxurious beach front resorts. This pension included spartan but clean bungalows amidst a horse ranch (the smell was definitely unique). The proprietor, Patrick, spoke English with a French accent but he claimed to be one of a long line of Spanish caballeros; he moved to Raiatea from

the Marquesas with 40 horses about 15 years earlier and set up a business as a guide (to the interior and the Temehanirahi summit plateau of Raiatea) as well as lead horse-back rides locally.

Ron immediately went to work setting up his mobile photo studio and once finished, we hopped in the Land Rover to scout out potential streams and waterfalls (cascades) to sample the next four days on the island. One particularly good one, the Mitimitiaute Riviere happened to be next door to the pension and had a good road to a trailhead that led to the “Trois Cascades”. At the junction of the trail and the stream, we did some preliminary sweeping of stream riffles and collected a few *Campsicnemus* swarming among the more dense simuliids (this turns out to be a typical behavior for them in FP). I set up some pan traps to be collected in a few days and after hiking up stream a bit, we called it a day.

Wednesday, we drove to the west side of the island, parked the truck at the beach (where I swept the beach rubble and got a few beach simuliids), and hiked a few kilometers up a soggy road through hibiscus, then *Casuarina*, and finally a muddy and slippery rain-eroded trail through *Metrosideros* up to where the vegetation opened up and the trail crossed a *Pandanus*-lined stream cascading over weathered sandstone bedrock (Fig. 3). The stream drained the summit (Temehanirahi) plateau (about 430 m).



Fig. 3. Stream on top of Temehanirahi plateau.

We sampled the stream and vegetation and set up pan traps and a Malaise trap. A few *Chrysotus* were collected at the stream and trap site -- the pan traps (and Malaise trap) were collected a few days later by Elin but did not have any *Campsicnemus* (although collecting the previous year at the summit a few hundred meters above had them). Previous rains in the area may have scoured out the stream area since there wasn't much flying that day although we had sunny weather. The hike up and down took much of the day, so there wasn't much daylight left to collect once we got back to the beach and the truck and headed back to our bungalow.

traditional dichotomous keys. We strongly encourage authors to include links to regional databases, maps, and related products, and we insist that all submissions to this series include high-quality illustrations and/or photographs (see instructions to authors on the website for technical recommendations regarding format, image size, etc).

Editorial board, Canadian Journal of Arthropod Identification:

S.A. Marshall, Guelph (editor)
 V.M. Behan-Pelletier, Ottawa (Soil arthropods)
 C.M. Buddle, Montreal (Other terrestrial arthropods)
 J.M. Cumming, Ottawa (Diptera – Lower Brachycera)
 D.C. Currie, Toronto (Diptera – Nematocera)
 H.V. Danks, Ottawa (BSC secretariat)
 T.D. Galloway, Winnipeg (Parasitic Insects)
 D.J. Giberson, Charlottetown (Aquatic Insects)
 J.H. Huber, Ottawa (Hymenoptera)
 J.F. Landry, Ottawa (Lepidoptera)
 R.E. Roughley, Winnipeg (Coleoptera)
 G.G.E. Scudder, Vancouver (Hemipteroids)
 F.A.H. Sperling, Edmonton (Lepidoptera)
 T.A. Wheeler, Montreal (Diptera – Higher Brachycera)

Good News! Re: Manual of Central American Diptera

The editorial board has been hard at work this winter and springtime on the completion of the Manual. Final stages of editing are nearly finished and authors will be receiving a version of their chapter(s) at the beginning of May for final, minor, adjustments. Volume 1 will be submitted to the National Research Council of Canada in May for publication and it is expected that Volume 2 will be submitted by this fall. At the present time we are uncertain as to what the volumes will cost but will keep you informed. The end is near!

Awesome Book

by Art Borkent
 691-8th Ave. SE, Salmon Arm, British Columbia,
 V1E 2C2, Canada; aborkent@jetstream.net

I just finished reading “Endless Forms Most Beautiful” by Sean B. Carroll (2005) and want to recommend this book to all Dipterists. It provides an overview of the connection between genes and the form of organisms, explaining how genes end up providing the structure and organization of the organisms many of us study. The processes of development, including the underlying organizational processes that produce a complex organism, are explained in plain language, with plenty of examples

and illustrations, that is a pleasure to read. I recommend it to anyone who wants to better understand the area of evolutionary development (don't we all?). It certainly helped me to think more deeply about how gene sequences are translated into the synapomorphies I use to build my phylogenies of Ceratopogonidae and other Diptera.

Books and Publications

(with thanks to Chris Borkent, McGill, Montreal, Quebec, for completing a literature search)

- Brooks, S.E., J.M. Cumming, J.E. O'Hara, J.H. Skevington and B.E. Cooper. 2007. Diptera types in the Canadian National Collection of Insects. Supplement. Third edition. North American Dipterists Society Website [<http://www.nadsdiptera.org/catalogs/CNCtypes/suppl.htm>].
- Cameron, S. L., C.L. Lambkin, S.C. Barker, and W.F. Whiting. 2007. A mitochondrial genome phylogeny of Diptera: whole genome sequence data accurately resolve relationships over broad timescales with high precision. *Systematic Entomology*. 32:40-59.
- Dittmar, -K., M.L. Porter, S. Murray, and M.F. Whiting. 2006. Molecular phylogenetic analysis of nycteribiid and streblid bat flies (Diptera: Brachycera, Calyptratae): Implications for host associations and phylogeographic origins. *Molecular Phylogenetics and Evolution* 38:155-170.
- Green, A.J. and M.I. Sánchez. 2006. Passive internal dispersal of insect larvae by migratory birds. *Biology Letters* 2:55-57. [description of live chironomid larvae being passed by black-tailed Godwits]
- Lonsdale, O, and S.A. Marshall. 2006. Redefinition of the Clusiinae and Clusiodinae, description of the new subfamily Sobarocephalinae, revision of the genus *Chaetoclusia* and a description of *Procerosoma* gen. n. (Diptera : Clusiidae). *European Journal of Entomology*. 103:163-182.
- Roháček, J. 2006. A monograph of Palaearctic Anthomyzidae (Diptera), Part 1. 16.5 x 24 cm, 326 pages, 661 black-and-white illustrations. Published as supplement 1 of the *Časopis Slezského zemského muzea*, Vol. 55. ISSN 1211-3026, ISBN 80-86224-57-0. Price (including postage and bank transfer taxes) 35 Euros. Orders should be sent to: Slezské zemské muzeum, Tyršova 1, 746 01 Opava, Czech Republic or by e-mail to mikulcova@szmo.cz, or david.fiser@szmo.cz
- All the Palaearctic taxa of Anthomyzidae are monographed. In the first part of the monograph, the history of taxonomic investigations on the Palaearctic Anthomyzidae, the morphology of preimaginal stages (egg, larva, puparium) and adults, and the systematic position and higher classification of Anthomyzidae are reviewed. The systematic treatments of all taxa (including fossil ones) of Anthomyzidae recognized in the Palaearctic Region are presented, with most complete data dealing with nomenclature, type material, taxonomy (diagnoses and redescriptions, keys, illustrations of taxonomically important structures and diagnostic features), preimaginal stages, relationships, biology and distribution. Two subfamilies, Protanthomyzinae (with one fossil genus and one species) and Anthomyzinae (with 11 extant genera and 42 species), are recognized in the region. Keys to the identification of all recognised taxa are presented.

- Stevens, J.R. and J.F. Wallman. 2006. The evolution of myiasis in humans and other animals in the Old and New Worlds (part I): phylogenetic analyses. *Trends in Parasitology*. 22:129-136.
- Stevens, J.R. and J.F. Wallman, D. Otranto, R. Wall, and T. Pape. 2006. The evolution of myiasis in humans and other animals in the Old and New Worlds (part II): biological and life-history studies. *Trends in Parasitology* 22:181-188.
- Sultana, F., Y.-G. Hu, M.J. Toda, K. Takenaka, M. Yafuso. 2006. Phylogeny and classification of *Colocasiomyia* (Diptera, Drosophilidae), and its evolution of pollination mutualism with aroid plants. *Systematic Entomology* 31:684-702.

Submission Form for Directory of North American Dipterists

For those who have not yet sent in a synopsis of their interests for the *Directory of North American Dipterists*, the following form is provided. Please restrict yourselves to no more than 20 words when listing the titles of your major projects and the animals you work with. Should any of you like to expand or modify your entries from the last list, use the form to indicate the changes.

The information can be emailed, or the form completed and faxed or sent to the following address:

Dr. J. M. Cumming,
Invertebrate Biodiversity
Agriculture & Agri-Food Canada,
K.W. Neatby Building, C.E.F.
Ottawa, Ontario, CANADA, K1A 0C6

FAX: (613) 759-1927

Email: cummingjm@agr.gc.ca

Full name: _____ **Address:** _____

_____ **Telephone Number:** _____

FAX Number: _____ **Email:** _____

Projects and taxa studied: _____
